

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.


In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

## Landscape and Urban Planning

journal homepage: [www.elsevier.com/locate/landurbplan](http://www.elsevier.com/locate/landurbplan)

# Estimating the removal of atmospheric particulate pollution by the urban tree canopy of London, under current and future environments

Matthew Tallis<sup>a</sup>, Gail Taylor<sup>a</sup>, Danielle Sinnott<sup>b,1</sup>, Peter Freer-Smith<sup>b,\*</sup>

<sup>a</sup> School of Biological Sciences, Life Sciences Building, University of Southampton, SO17 1BJ, UK

<sup>b</sup> Forest Research, Alice Holt Lodge, Farnham, Surrey GU10 4LH, UK

## ARTICLE INFO

## Article history:

Received 1 February 2011

Received in revised form 6 July 2011

Accepted 7 July 2011

Available online 1 September 2011

## Keywords:

Climate change

Greater London Authority

PM<sub>10</sub>

UFORE

Air quality

## ABSTRACT

Urban green space and particularly the tree canopy have been highlighted as offering a mitigation potential against atmospheric particulate pollution. In this paper current and future particulate (PM<sub>10</sub>) deposition to the urban tree canopy of the Greater London Authority (GLA) was estimated. A modelling approach was used based on the Urban Forest Effects Model (UFORE) and a modified version. Here we give evidence showing that these deposition models can be adapted to run from annual mean meteorological and PM<sub>10</sub> concentration data, thus providing a methodology to examine future scenarios.

Depending on the modelling approach, the urban canopy of the GLA is currently estimated to remove between 852 and 2121 tonnes of PM<sub>10</sub> annually; representing between 0.7% and 1.4% of PM<sub>10</sub> from the urban boundary layer. Estimates of PM<sub>10</sub> removal which take into account a planned increase in tree cover, from the current 20% to 30% of the GLA land area, suggest deposition of 1109–2379 tonnes (1.1–2.6% removal) by the year 2050. The evidence provided here suggests that the targeting of tree planting in the most polluted areas of the GLA and particularly the use of street trees which have the greatest exposure to PM<sub>10</sub>, would have the greatest benefit to future air quality. The increased deposition would be greatest if a larger proportion of coniferous to broadleaved trees were used at such sites.

© 2011 Elsevier B.V. All rights reserved.

## 1. Introduction

In the urban environment reductions in air quality resulting from emissions of particulate matter, primarily from road traffic, are a serious health issue globally (Gupta, Kumar, Maharaj Kumari, & Srivastava, 2004; Nowak, Crane, & Stevens, 2006; Yang, McBride, Zhou, & Sun, 2005). Estimates for the UK indicate that short-term exposure to the levels of particulate matter of less than  $10 \times 10^{-6}$  m in aerodynamic diameter (PM<sub>10</sub>) led to an additional 6500 deaths and 6400 hospital admissions in 2002 (AQEG, 2005). The UK Department of Health also estimates that 1.9% of urban deaths can be attributed to PM<sub>10</sub> pollution (COMEAP, 1998). Vegetation captures gases, particulates and aerosols from the atmosphere more effectively than other land surfaces (Fowler, Cape, & Unsworth, 1989; Smith, 1981). As a result of their large canopy surface area of leaves, stem and branches and the air turbulence created by their

structure, trees are more effective in the uptake of these materials than shorter vegetation (Lovett, 1994; Powe & Willis, 2004). For example, Fowler et al. (2004) found that woodlands in the West Midlands of England collect three times more PM<sub>10</sub> than grassland. Brownian motion accounts for the flux of gases and fine particles along concentration gradients (diffusion) and this mechanism is important for pollutants that dissolve on wet leaf surfaces. Larger tree leaf areas will thus increase the uptake of fine particles by diffusion. However, turbulent air flow and associated impaction are the main mechanisms resulting in the greater deposition of particles to trees as compared to that to shorter vegetation. The inertia of particles travelling in an air stream as it curves around an object, such as a leaf or stem, forces them through the boundary layer and onto the object. Thus the canopy area and structure (i.e. tree species), concentration of particles in the airstream, particle size distribution, and windspeed are all important factors in determining particle uptake by vegetation (see a full discussion in Beckett, Freer-Smith, & Taylor, 2000c), and these factors are the main inputs required to model particle deposition (see Section 2.1). Two deposition models, the Urban Forest Effects Model (UFORE; Nowak & Crane, 1998) and FRAMES (Bealey et al., 2007), have been developed to evaluate PM<sub>10</sub> uptake by trees in urban areas. Recently Tiwary et al. (2009) used a pollution/deposition flux approach and species-specific deposition velocities (Freer-Smith, Beckett, & Taylor, 2005; Freer-Smith,

\* Corresponding author. Tel.: +44 01420 526211.

E-mail addresses: [m.j.tallis@soton.ac.uk](mailto:m.j.tallis@soton.ac.uk) (M. Tallis), [g.taylor@soton.ac.uk](mailto:g.taylor@soton.ac.uk) (G. Taylor), [danielle.sinnott@uwe.ac.uk](mailto:danielle.sinnott@uwe.ac.uk) (D. Sinnott), [peter.freer-smith@forestry.gsi.gov.uk](mailto:peter.freer-smith@forestry.gsi.gov.uk) (P. Freer-Smith).

<sup>1</sup> Current address: Centre for Sustainable Planning and Environments, Department of Planning and Architecture, University of the West of England, Frenchay Campus, Coldharbour Lane, Bristol BS16 1QY, UK.

El-Khatib, & Taylor, 2004) to estimate PM<sub>10</sub> capture in a case study area of London, UK.

There is a need for urban greenspace to be established strategically at a landscape scale in order to maximise the full range of potential benefits it can bring. These benefits include improving air quality, urban drainage, aesthetics, urban biodiversity, climate amelioration, enhancing recreation, conservation and health and well-being. The Read report suggests these services will help urban populations adapt to climate change (Read et al., 2009). Using a 10,000 ha study area within the Greater London Authority (GLA) (estimated as 6.4% of GLA land area) Tiwary et al. (2009) suggests that the implementation of the East London Green Grid with 25% tree cover could result in the aversion of two deaths and two hospital admissions per year through PM<sub>10</sub> removal. While Powe and Willis (2004) suggest that current woodland cover in Great Britain mitigates between five and seven deaths and four and seven hospital admissions annually, through PM<sub>10</sub> and SO<sub>2</sub> removal.

Dry deposition of PM<sub>10</sub> to trees primarily results from sedimentation under gravity and impaction under the influence of wind (Nowak, 1994a) and is therefore less influenced by rapidly fluctuating meteorology, but is largely affected by leaved canopy duration and total canopy leaf area (Nowak et al., 2006). During the hot summers of 2003 and 2006 experienced in the UK urban PM<sub>10</sub> concentrations, mostly the <PM<sub>2.5</sub> size fraction, increased through enhanced photochemical formation of secondary particles (Bower et al., 2008). Future temperatures are predicted to increase, particularly in urban areas, so understanding the role of trees is needed to maximise future benefits to urban air quality. The modelling approach used in this study has allowed an estimation of the role of urban trees in removing PM<sub>10</sub> from urban air under current and future environments.

Here we describe a detailed analysis of PM<sub>10</sub> uptake by tree from dry deposition in the c. 157K ha area covered by the Greater London Authority (GLA see Fig. 1); one of five European cities selected as case studies in the EU FP 7 Project “Sustainable Urban planning Decision support accounting for metabolism” (BRIDGE; [www.bridge-fp7.eu](http://www.bridge-fp7.eu)). In order to determine the relationships between amount and type of tree cover and PM<sub>10</sub> uptake we have used two modelling approaches: (i) the pollution/deposition flux method which has been used in a number of studies (e.g. Lovett, 1994; Nowak, 1994a; Nowak & Crane, 1998; Smith, 1981) and (ii) the approach used by Tiwary et al. (2009) where species specific deposition velocities were used with the pollution flux approach – referred to here as “the Tiwary method”. We have used these methods to estimate the PM<sub>10</sub> deposition to street trees, urban woodland, garden and other trees (remainder) for the whole GLA for current and future (2050) PM<sub>10</sub> capture. We also set out the methodology which allows these estimates to be made from annual input data of PM<sub>10</sub> concentrations and meteorology. This is an approach which can be taken for other cities where similar data are available.

## 2. Methodological strategy

Typically assessments of the capture of PM<sub>10</sub> by the urban tree canopy are carried out as part of an assessment of the overall air pollution removal potential (e.g. Donovan, Stewart, Owen, MacKenzie, & Hewitt, 2005; Nowak et al., 2006; Yang et al., 2005). This consists of running uptake models using local hourly meteorological and pollution data (e.g. Nowak & Crane, 1998). Here we investigated the potential of using existing tree survey data and annual maps of PM<sub>10</sub> distribution and observed/predicted meteorological conditions to estimate current and future PM<sub>10</sub> removal by the whole urban canopy of the GLA.

### 2.1. The models

The downward PM<sub>10</sub> flux ( $F$ ; in  $\text{g m}^{-2} \text{s}^{-1}$ ) to the urban tree canopy can be calculated as the product of the deposition velocity ( $V_d$ ; in  $\text{m s}^{-1}$ ) and pollutant concentration ( $C$ ; in  $\text{g m}^{-3}$ ) according to the methodology used in a number of studies (e.g. Lovett, 1994; Nowak, 1994b; Smith, 1981).

$$F = V_d \cdot C \quad (1)$$

$V_d$  is calculated as the inverse of the sum of the aerodynamic ( $R_a$ ), quasi-laminar boundary layer ( $R_b$ ) and canopy ( $R_c$ ) resistances (Baldocchi, Hicks, & Camara, 1987).


$$V_d = (R_a + R_b + R_c)^{-1} \quad (2)$$

To give PM<sub>10</sub> flux (PM<sub>10</sub> capture) the deposition value is multiplied by the area of the surface (e.g. canopy covered land area for the pollution flux method or canopy covered land area multiplied by the Leaf Area Index (LAI) for the Tiwary method) over time periods for which the pollutant concentration is known e.g. hourly or annually. Leaf Area Index is the ratio of total tree leaf area to the projected ground area occupied by the canopy ( $\text{m}^2 \text{m}^{-2}$ ). Local meteorological conditions are used to calculate the resistances  $R_a$  and  $R_b$  in  $\text{s m}^{-1}$  as described in detail in Nowak, 1994b. Canopy resistance ( $R_c$ ;  $\text{s m}^{-1}$ ) values are derived from average  $R_a$  and  $R_b$  and deposition velocity ( $V_d$ ; in  $\text{m s}^{-1}$ ) as described by Nowak, 1994b and Tiwary et al. (2009). It is within the  $V_d$  parameter that the Tiwary method differs.

In the UFORE model a maximum value of  $V_d$  is set at  $0.0064 \text{ m s}^{-1}$  based on the median deposition velocity from the literature (Lovett, 1994) and a 50% re-suspension rate (Zinke, 1967) as described by Nowak and Crane (1998). This  $V_d$  assumes a single sided LAI of  $6 \text{ m}^2 \text{m}^{-2}$ , 6% coniferous cover and deposition to stems with a bark surface area of  $1.7 \text{ m}^2 \text{m}^{-2}$  of land area giving a total plant area index of  $7.7 \text{ m}^2 \text{m}^{-2}$ . Here we assigned a  $V_d$  of  $0.0064 \text{ m s}^{-1}$  (Summer),  $0.0039 \text{ m s}^{-1}$  (Spring and Autumn) and  $0.0014 \text{ m s}^{-1}$  (Winter) for broadleaved trees to represent a change in LAI from an urban peak of  $6 \text{ m}^2 \text{m}^{-2}$  to a Spring and Autumn mid value of  $3 \text{ m}^2 \text{m}^{-2}$  and also accounting for deposition to bark in Winter. We assigned the 4% coniferous component of the GLA an annual LAI of  $6 \text{ m}^2 \text{m}^{-2}$  and therefore an annual  $V_d$  of  $0.0064 \text{ m s}^{-1}$ ; this is a likely under-estimate of true urban coniferous canopy LAI but allows for a comparison of the pollution flux model approach to be made with results from the UFORE model applied to multiple USA cities in which a LAI of  $6 \text{ m}^2 \text{m}^{-2}$  is assumed (Nowak et al., 2006). As  $V_d$  includes an LAI and bark area parameter the PM<sub>10</sub> flux in Eq. (1) is multiplied by the land area to give total flux ( $F$ ). Annual stem deposition was calculated as total broadleaf Winter deposition multiplied by four (Spring, Summer, Autumn and Winter) and deposition to foliage was calculated from the difference between total canopy deposition and deposition to the stems.

Tiwary et al. (2009) used species-specific  $V_d$  values to the foliage of *Acer pseudoplatanus* to represent the broadleaved canopy (this species being the most common large broadleaf in the survey of Britt and Johnson, 2009) and *Pinus menziesii* representing coniferous canopy (which has similar  $V_d$  values to Leyland cypress, the most common conifer in the Britt and Johnson survey). Depositions were calculated using the known relationships between wind speed ( $3\text{--}9 \text{ m s}^{-1}$ ) and  $V_d$ , which have been established for *A. pseudoplatanus* and *P. menziesii* but not Leyland cypress (Freer-Smith et al., 2004). Here we apply these same relationships as given in Tiwary et al. (2009) to represent deposition to foliage of broadleaved and coniferous canopies. The annual mean wind speeds were  $3.6 \text{ m s}^{-1}$  and  $3.8 \text{ m s}^{-1}$  for 2004 and 2006 respectively as calculated from the seasonal maps (Perry & Hollis, 2005). The mean wind speed used in the current study was therefore within the ranges of wind speed under which measured data





**Fig. 1.** The Greater London Authority (GLA) displaying the administrative boundaries of the 33 boroughs and the mapped broadleaf and coniferous woodland cover. Also mapped are the locations of the four hectare sampling units used by Britt and Johnston (2008) to survey (■) garden and street tree (●) the location of the 11 LAQN background PM<sub>10</sub> monitoring sites (●) used for the background PM<sub>10</sub> seasonal correction factor and the location of figure two (■).

were collected (Freer-Smith et al., 2004). Using the mean wind speed for 2006 resulted in a deposition velocity of  $0.0021 \text{ m s}^{-1}$  and  $0.0109 \text{ m s}^{-1}$  to broadleaf and coniferous foliage respectively. Although the  $V_d$  were modelled with mean wind speeds, it is noted that windspeed will fall as air travels through a canopy so that estimates could be improved by more detailed modelling of urban airflow. As  $V_d$  is calculated to foliage area, PM<sub>10</sub> flux from Eq. (1) is multiplied by the land area and seasonal LAI to give total leaf flux. Similarly, stem deposition was modelled as a linear relationship with wind speed for *A. pseudoplatanus* and *P. menziesii* (Freer-Smith et al., 2004) and deposition velocities for the mean wind speed of  $3.8 \text{ m s}^{-1}$  were estimated. These relationships were  $Y = 0.2 \times 10^{-3} X$  ( $r^2 = 0.96$ ;  $n = 3$ ) and  $Y = 0.8 \times 10^{-5} X$  ( $r^2 = 0.83$ ;  $n = 3$ ), giving  $V_d$  values of  $7.6 \times 10^{-4} \text{ m s}^{-1}$  and  $27.2 \times 10^{-6} \text{ m s}^{-1}$  for stems of *A. Pseudoplatanus* and *P. menziesii* respectively. Given the surface area index for bark of  $1.7 \text{ m}^2 \text{ m}^{-2}$  this results in a  $V_d$  term to the land surface under broadleaved trees of  $0.0013 \text{ m s}^{-1}$  and  $0.000046 \text{ m s}^{-1}$  under coniferous. As deposition velocity to broadleaf stems was very similar to the UFORE value of  $0.0014 \text{ m s}^{-1}$ , and deposition to coniferous stems was lower (combined with a low area, was assumed negligible) then the UFORE deposition to stems was added to the leaf deposition here to give total deposition for the Tiwary method. Particles deposited to tree canopies either remain on the leaf and bark surfaces, are resuspended to the atmosphere or are

washed of by precipitation. Unlike UFORE, the Tiwary method assumes 0% re-suspension because there is now published experimental evidence that that resuspension is extremely small to tree canopies at wind speeds similar to those experienced in London (see Tiwary et al., 2009). In addition the  $V_d$  values used in the Tiwary method are based on measurements made in windtunnels and are calculated from net uptake; that is the final uptake after resuspension has occurred.

The percentage reduction in PM<sub>10</sub> within the whole mixing layer of the GLA caused by the urban canopy was estimated assuming a well mixed boundary layer which is common during daytime, stable conditions (Colbeck & Hanison, 1985). The total amount of PM<sub>10</sub> removed by the canopy over the year was expressed as a percentage of the total amount of PM<sub>10</sub> present in the mixing layer; the annual exposure where:

$$\text{Annual exposure} = \text{PM}_{10} (\mu\text{g m}^{-3}) \times \text{MLH (m)} \times 8760 \\ (\text{hours per year}) \times \text{LA (m}^2)$$

when PM<sub>10</sub> = mean annual PM<sub>10</sub> concentration, MLH = mean annual mixing layer height and LA = land area cover by urban canopy.

Annual PM<sub>10</sub> removal and mixing layer height were estimated using the seasonal models for current and future scenarios, annual mean PM<sub>10</sub> concentration were estimated from LAEI (2006) for the

**Table 1**  
Current and future estimates of urban tree canopy cover for the Great London Authority (GLA) reported as total cover and cover of each of the four canopy types. For 2050 two scenarios are presented; in 1 tree cover is increased across all woodland types to give 30% tree cover of the GLA land area and in scenario 2 cover is increased in only one woodland type – a to d to achieve 30% tree cover. Changes from the current tree cover are shown in bold.

Scenario	Canopy type				
	Total	Woodland	Street	Garden	Remainder
<i>Current tree cover</i>					
As a % of the total GLA	20	8.64	1.1	4.73	5.40
Total hectares	31,265	13,503	1782	7394	8586
<i>2050 scenario 1</i>					
As a % of the total GLA	<b>30</b>	<b>12.9</b>	<b>1.8</b>	<b>7.2</b>	<b>8.1</b>
Total hectares	<b>46,898</b>	<b>20,166</b>	<b>2814</b>	<b>11,256</b>	<b>12,662</b>
<i>2050 scenario S2a</i>					
As a % of the total GLA	<b>30</b>	<b>18.6</b>	1.1	4.73	5.4
Total hectares	<b>46,898</b>	<b>29,135</b>	1782	7394	8586
<i>2050 scenario S2b</i>					
As a % of the total GLA	<b>30</b>	8.64	<b>11.1</b>	4.73	5.40
Total hectares	<b>46,898</b>	13,503	<b>17,415</b>	7394	8586
<i>2050 scenario S2c</i>					
As a % of the total GLA	<b>30</b>	8.64	1.1	<b>14.8</b>	5.4
Total hectares	<b>46,898</b>	13,503	1782	<b>23,027</b>	8586
<i>2050 scenario S2d</i>					
As a % of the total GLA	<b>30</b>	8.64	1.1	4.73	<b>15.5</b>
Total hectares	<b>46,898</b>	13,503	1782	7394	<b>24,219</b>

current scenario and from Williams (2007) for the future scenario. Mixing layer height (MLH) was determined as follows:

$$MLH = \frac{275 \cdot c \cdot u_z}{(\sin(\varphi) \cdot \log(z - d/z_0))}$$

where  $\varphi$  is latitude,  $c$  is a parameter related to stability classes using the stability classification scheme of Pasquill,  $u_z$  is the windspeed at canopy height ( $z$ ),  $d$  is the displacement height ( $d = 0.7z$ ) and  $z_0$  is the aerodynamic roughness length ( $z_0 = 0.1z$ ) and  $z$  = canopy height (10 m); see detail and references in Tiwary et al. (2009).

## 2.2. Quantifying the urban tree canopy for London

A combination of published data from ground, airborne and satellite surveys were used to classify and quantify the tree canopy cover for London. The GLA borough boundaries and water features were visualised from the 1:20,000 scale vector boundary data maps (Great Britain boundary data, Collins Bartholomew, UK) using Geographic Information Systems (GIS) software (ERSI® ArcMap version 9.3, ESRI, Aylesbury, UK) with the OSGB\_1936 coordinate system and the Transverse Mercator projection.

For each of the 33 London boroughs we were able to quantify canopy cover as (i) urban woodland, (ii) street trees, (iii) garden trees and, for the whole GLA area, data were also available for a fourth class (iv) remainder (Fig. 1). Urban woodland was mapped to the GLA boundary using the 25 m<sup>2</sup> resolution land classification map (Fuller, Smith, Sanderson, Hill, & Thompson, 2001). This provides the spatially referenced mapped woodland data as can be seen in Fig. 1. The 25 m raster data were extracted for land cover classes 1.1 (broadleaved woodland) and 2.1 (coniferous woodland). The broadleaved class comprised broadleaved woodlands of stands >5 m high with tree cover of >20%; scrub (<5 m) requires cover >30% for inclusion. The coniferous canopy represents only c. 4% of the urban tree cover of the GLA area and includes stands and plantations of >20% cover, more details can be found in Fuller et al. (2002). We took the borough land area and the estimated canopy cover of street trees for ten London boroughs from Britt and Johnston (2008) to estimate street tree area for each borough giving 1.1% cover of land area. Additionally, the area of land covered by street trees in each borough was estimated by multiplying the number of street trees in each borough (GLA, 2007) with the mean estimated ground area under street tree canopy (Britt & Johnston, 2008) giving 0.94% coverage of land area. The coverage of trees in

residential gardens was also estimated from Britt and Johnston (2008) and the locations of the sample survey sites used by these authors to quantify the land area occupied by street and garden canopy cover are shown in Fig. 1. Table 1 shows the breakdown of canopy classes discussed. Urban woodland, street and garden trees represent 22,286 ha which is 72.5% of the total reported canopy area. The further 27.46% (8441 ha) was classed as 'remainder'. This class could include individual trees in parks and other green spaces, woodland of less than 25 m<sup>2</sup>, and street or garden trees missed due to the spatially limited sampling areas of these two canopy types.

For trees which are not spatially referenced canopy area can only be localised to the borough or whole GLA, and therefore model input data of PM<sub>10</sub> concentrations and meteorology had to be obtained that represent the whole GLA. Spatially averaged PM<sub>10</sub> concentrations and meteorology values for the whole GLA area were calculated from mapped PM<sub>10</sub> dispersion data that exists as annual mean values and those for 2006 were used (LAEI, 2006) and monthly mean mapped meteorological data (Perry & Hollis, 2005). The ability to use such input data would also allow the potential benefits of future tree planting to be modelled using future predictions of climate and PM<sub>10</sub> concentrations, which are typically estimated as annual means. In order to estimate the effects of using annual rather than hourly inputs we used both the pollution/deposition flux approach and the Tiwary method with hourly and seasonal input data and have compared the annual mean resistance parameters and canopy PM<sub>10</sub> uptake values.

## 2.3. Future canopy cover

The GLA has announced the ambitious aim of increasing urban tree cover from the current area (2006–08) of 20% of the whole GLA land surface area to 30% by 2050 (GLA, 2009), which could be achieved through the planting of trees on land managed by the boroughs or through encouraging tree planting on private land. To determine the influence of this increase on PM<sub>10</sub> removal from the atmosphere we ran models for five future scenarios of tree cover (Table 1). In the first (S1), the increase in tree cover was obtained by increasing each canopy type (i.e. urban woodland, street trees, garden trees and remainder) so they contributed the same proportional cover as they do currently. Secondly in scenarios S2a to S2d, the increase was achieved by increasing only one of each of the canopy types rather than all four. Thus under scenario S2a urban

woodland was increased by increasing the areas of street trees, with the area of garden trees and remainder left unaltered. In S2b street tree cover was increased, in S2c garden trees and S2d the cover of remainder trees was increased. The current distribution of 96% broadleaf and 4% coniferous was used in all the scenarios examined and we have assumed that the newly planted areas will reach normal canopy size by 2050. This is a reasonable assumption given the time frame and the very common use of large containerized trees in urban plantings.

#### 2.4. The seasonal adaptations

To estimate the current PM<sub>10</sub> removal by the urban canopy the models were run using both hourly input and annual input values using both the pollution flux method and the Tiwary method. This allowed us to examine the difference in the outputs from these two approaches and to identify the extent to which seasonal values derived from annual means can be used where hourly data are not available. For the GLA annual mean data were available for future predictions of meteorology (UKCP09) and for PM<sub>10</sub> concentrations (Williams, 2007). In order to take account of seasonal changes in canopy LAI we used different values for the LAI for winter, spring, summer and autumn. The LAI for broadleaf trees and coniferous trees was set as described earlier and season start and end dates are those used by UKCP09. Daily mean PM<sub>10</sub> concentration data for 2006 were downloaded from 11 Local Air Quality Network monitoring sites (LAQN) in order to provide background PM<sub>10</sub> concentrations ([www.londonair.org.uk/](http://www.londonair.org.uk/)). These 11 sites were those that remained after rejection of those where site-specific factors gave peak values. These monitoring data were used to calculate seasonal correction factors from the fraction of the annual mean which was represented by each seasonal mean. An identical approach was taken using monitoring stations designated as 'street' ( $n = 21$  stations) and the same mean seasonal correction factor values resulted. Annual background and street PM<sub>10</sub> concentrations were then converted to seasonal values using this factor.

#### 2.5. Meteorological and PM<sub>10</sub> concentration input data

##### 2.5.1. Current

Hourly meteorological data for 2004 from Heathrow airport ( $-0^{\circ}44.9'W$ ,  $51^{\circ}47.9'N$ ) was obtained from the Meteorological Office (Meteorological Office, 2006). These data were used in both the model runs using hourly inputs and to calculate the seasonal means for the seasonal inputs. The mapped seasonal data for the GLA were extracted from the UKCP09 gridded data sets of monthly values ([www.metoffice.gov.uk](http://www.metoffice.gov.uk); Perry & Hollis, 2005) for 2004 and 2006 at a 5 km resolution for the whole GLA area. Seasonal mean data were calculated using the mosaic function in ArcMap® and designated as 'seasonal' from mapped model inputs.

To represent the mean hourly GLA background concentrations, hourly background PM<sub>10</sub> concentrations were obtained from the Islington 1 LAQN site (the mean daily data at this site closely represent the mean daily background concentrations of the 11 sites selected for the background correction factor). A seasonal average PM<sub>10</sub> concentration was also calculated from this data (seasonal from hourly means). Seasonal PM<sub>10</sub> concentration data were also calculated from PM<sub>10</sub> distribution maps for the GLA (LAEI, 2006) for 2006 (seasonal from maps). Seasonal street level PM<sub>10</sub> concentrations were obtained from a 20 m buffer applied to the GLA road network using ArcMap® (Fig. 2). The mean 20 m street buffer PM<sub>10</sub> concentration for the whole GLA was  $23.5 \mu\text{g m}^{-3}$  (mean borough range: Bromley  $21.7 \mu\text{g m}^{-3}$  to City  $26.8 \mu\text{g m}^{-3}$ ) and the remaining background was  $21.7 \mu\text{g m}^{-3}$  (range: Bromley  $20.5 \mu\text{g m}^{-3}$  to City  $24.7 \mu\text{g m}^{-3}$ ). A large spatial concentration range within the

street tree buffer was evident (as exemplified by Fig. 2) ranging from  $19.9 \mu\text{g m}^{-3}$  to a GLA maximum of  $83.4 \mu\text{g m}^{-3}$  from the 20 m street buffer for 2006. To estimate the current capture of PM<sub>10</sub> by the urban woodland canopy, models were run using the hourly input data and the seasonal data (calculated as described above). This allowed for an analysis of the effect of using hourly and seasonal input data for meteorology and PM<sub>10</sub> exposure, the results are given in Tables 2 and 3.

#### 2.5.2. Future estimates

Input data to describe the future climate in the GLA were extracted from UKCP09 (<http://ukclimateprojections-ui.defra.gov.uk>). The absolute values (for a 50% probability) were extracted from the 'UK probabilistic projections of climate change over land' based on the medium emissions scenario (SRES A1B) for London in the 2050s. Seasonal change in wind speed for the London area was estimated from the UKCP02 maps for the mean of the medium emissions scenarios for 2080 (the only data available at the time of analysis). These estimates for wind speed are increases of 6%, 1.5% and 1.5% for winter, spring and summer respectively and a decrease of 4% for the autumn. The annual mean PM<sub>10</sub> concentrations in London for the year 2050 were taken from Williams (2007) who estimated as  $17.5$  and  $21 \mu\text{g m}^{-3}$  for background and roadside respectively. In calculating PM<sub>10</sub> concentrations the same seasonal adjustment factors were used for 2006 and 2050.

### 3. Results

#### 3.1. Current and future urban canopy cover

The current and future scenarios (S1 and S2a-d) of urban canopy cover for the GLA are given in Table 1 (urban woodland, garden trees, street trees and remainder). Both the available estimates of street tree canopy area are close at 1.1% (estimated from Britt & Johnston, 2008 area survey) and 0.94% (estimated from Britt & Johnston, 2008 canopy spread class survey and borough tree inventories, GLA, 2007). By neglecting the remainder class (which cannot be resolved at borough level), canopy cover per borough ranged from 5.9% (City) to 23.7% (Bromley). Fig. 1 clearly demonstrates the lack of urban forest included in the survey of Britt and Johnston (2008) which estimated the GLA to have only 8.2% tree canopy cover.




Fig. 2. An example area of the PM<sub>10</sub> concentration map (LAEI, 2006) showing distribution of PM<sub>10</sub> within the 20 m road buffer used to extract the street level PM<sub>10</sub> concentrations.



**Table 2**

The effect of changing input data from hourly to seasonal on the estimated values of mixing layer height, frictional velocity and the resistance parameter ( $R_a$ ,  $R_b$  and  $R_c$ ) for both the pollution/deposition and Tiwary approaches. Seasonal  $R_c$  values are given for the pollution flux method as the model estimates deposition to a canopy covered land surface with seasonal LAI. The Tiwary method estimates deposition to foliage therefore  $R_c$  remains a single value through the season but LAI changes.

Model inputs [PM <sub>10</sub> ] and meteorological data	Mixing height (m)	Frictional velocity (m s <sup>-1</sup> )	$R_a$ (s m <sup>-1</sup> )	$R_b$ (s m <sup>-1</sup> )	$R_c$ (s m <sup>-1</sup> )				
					Poll./Dep. method assuming a 10% coniferous canopy			Tiwary approach	
					Winter	Spring + Autumn	Summer	Broadleaf On leaf	Coniferous On leaf
Hourly	319.8	4.36	0.31	0.13	713.85	255.00	155.81	471.6	103.7
Seasonal calculated from hourly	388.4 <sup>a</sup>	4.45	0.22	0.04	714.03	256.15	156.00	447.5	82.4
Seasonal calculated from hourly and map	409.0	3.82	0.25	0.05	713.98	256.11	155.95	484.2	98.4
Seasonal from Map	409.0	3.82	0.25	0.05	713.98	256.11	155.95	484.2	98.4

<sup>a</sup> The mean of seasonal day and night estimates.

**Table 3**

The effect of changing input data from hourly to seasonal on the estimated PM<sub>10</sub> capture of the urban woodland component (foliage and stems, model outputs) of the total GLA urban canopy using the flux/deposition model approach.


Model inputs [PM <sub>10</sub> ] and meteorological data	Annual PM <sub>10</sub> deposition to urban woodland (tonnes)			Annual PM <sub>10</sub> deposition to urban woodland (kg ha <sup>-1</sup> y <sup>-1</sup> )	
	Broadleaf (sum of all seasons)	Coniferous	Total	Broadleaf	Coniferous
Hourly	327.7	22.1	348.8	25.3	41.8
Seasonal calculated from hourly	337.4	22.9	360.3	26.0	43.4
Seasonal from hourly and map	337.4	22.9	360.3	26.0	43.4
Seasonal from Map	344.8	23.1	367.9	26.6	43.8

### 3.2. Testing the seasonal model

The mixing height, frictional velocity and resistance parameters  $R_a$ ,  $R_b$  and  $R_c$  (aerodynamic, quasilinear boundary layer and canopy resistance terms respectively) are given in Table 2. The values given are for the pollution/deposition and Tiwary models using hourly or seasonal input data. All terms showed small changes as a result of scaling from hourly to seasonal inputs and the canopy terms ( $R_c$ ) are expressed differently because of adaptations made by Tiwary et al. (2009). The total PM<sub>10</sub> capture in 2004 by the urban woodland component of the GLA's canopy cover are shown in Table 3 for the pollution flux approach and in Table 4 for the Tiwary method. Estimates of PM<sub>10</sub> capture are given based on hourly and seasonal input parameters. For the pollution/deposition flux method the total annual PM<sub>10</sub> deposition ranged from 348.8 tonnes per year (t y<sup>-1</sup>) using hourly data to 367.9 t y<sup>-1</sup> using seasonal inputs; a difference of 19 t y<sup>-1</sup> (only 4%) (Table 3). Using this approach deposition was 27 kg ha<sup>-1</sup> y<sup>-1</sup> and 44 kg ha<sup>-1</sup> y<sup>-1</sup> to broadleaved and coniferous canopies respectively with seasonal input data and was very similar when hourly data were used (25 kg ha<sup>-1</sup> y<sup>-1</sup> for broadleaves and 42 kg ha<sup>-1</sup> y<sup>-1</sup> for conifers). With the Tiwary method deposition values are greater ranging from 985 t y<sup>-1</sup> when based on hourly data to 756 t y<sup>-1</sup> with seasonal inputs; a difference of 229 t y<sup>-1</sup> (30%) (Table 4). Deposition to conifers (between 422 kg ha<sup>-1</sup> y<sup>-1</sup> and 709 kg ha<sup>-1</sup> y<sup>-1</sup>) was considerably greater than to broadleaves (41 kg ha<sup>-1</sup> y<sup>-1</sup> to 47 kg ha<sup>-1</sup> y<sup>-1</sup>). Estimates of mean annual mixing layer height were 320 m using hourly input data and 410 m using seasonal inputs, Nowak and Crane (1998) estimate a minimum urban boundary layer height of 150 m and 250 m for night and day respectively.

### 3.3. PM<sub>10</sub> removal by the current and future (2050) GLA tree cover

Total annual estimates of PM<sub>10</sub> removal by dry deposition to the urban canopy (assuming a 96% broadleaf and 4% coniferous distribution) are given in Fig. 3. Across all scenarios and for 2006 and 2050 the Tiwary method gives an approximate 2.5 fold increase



**Fig. 3.** The estimates of annual PM<sub>10</sub> capture by the urban canopy of the GLA using (a) the pollution/deposition and (b) Tiwary models. Values are given for direct model outputs; therefore values are for deposition to foliage and stems for the deposition approach and foliage alone for the Tiwary method. Current canopy cover in 2006 meteorology and PM<sub>10</sub> environment, current cover in 2050 meteorology and PM<sub>10</sub> environment and four different future scenarios of canopy cover in 2050 meteorology and PM<sub>10</sub> environment are given.

**Table 4**

The effect of changing input data from hourly to seasonal input data on the estimated PM<sub>10</sub> capture of the urban woodland component (foliage only, model outputs) of the total urban canopy using the Tiwary approach.

Model inputs	Annual PM <sub>10</sub> deposition to urban woodland foliage (tonnes)			Annual PM <sub>10</sub> deposition to urban woodland foliage (kg ha <sup>-1</sup> y <sup>-1</sup> )	
	Broadleaf	Coniferous	Total	Broadleaf	Coniferous
[PM <sub>10</sub> ] and meteorological data					
Hourly	609.6	375.1	984.7	47.0	709.8
Seasonal calculated from hourly	569.6	263.5	833.1	43.9	498.6
Seasonal from hourly and map	518.1	222.9	741.0	39.9	421.8
Seasonal from map	532.9	223.2	756.1	41.1	422.4

**Table 5**

The deposition of PM<sub>10</sub> to the urban canopy (kg ha<sup>-1</sup> y<sup>-1</sup>) estimated using both modelling approaches and with current canopy cover and 2006 meteorology and [PM<sub>10</sub>] taken from spatial maps of the GLA.

Canopy type	Flux/deposition model estimates of PM <sub>10</sub> capture (kg ha <sup>-1</sup> y <sup>-1</sup> )			Tiwary approach estimates of PM <sub>10</sub> capture (kg ha <sup>-1</sup> y <sup>-1</sup> )		
	GLA configuration (96%:4% broadleaf:coniferous)	100% broadleaf	100% coniferous	GLA configuration (96%:4% broadleaf:coniferous)	100% broadleaf	100% coniferous
Woodland, Garden and Remainder	27.3	26.6	43.8	58.5	42.2	448.2
Street trees	29.6	28.8	47.5	63.4	45.8	486.2

over the values derived from the original pollution flux method. This reflects the larger  $V_d$  values used in the Tiwary method, particularly for conifers at high wind speeds and also the 50% re-suspension rate included in the pollution flux method is not in the Tiwary approach. Total capture for 2006 is estimated at 852 tonnes, from the model output (559 tonnes to foliage) and 2121 tonnes (1828 tonnes to foliage) for the pollution flux and Tiwary approaches respectively. Table 5 gives the uptake of PM<sub>10</sub> for both these models in kg ha<sup>-1</sup> y<sup>-1</sup>. In both approaches uptake to street trees was slightly larger than to other canopy types and the Tiwary method gave a 10-fold increase of deposition to the coniferous canopy when compared to the pollution flux method. In the absence of *in situ* validation data, we compared our modelled estimates of PM<sub>10</sub> deposition to foliage with measured data of PM<sub>10</sub> deposition to urban foliage within the UK. The mean street level PM<sub>10</sub> concentration for the GLA was 23 µg m<sup>-3</sup> and the flux/deposition model estimates a PM<sub>10</sub> deposition of 638 mg m<sup>-2</sup> to foliage and the Tiwary approach 1533 mg m<sup>-2</sup>. Deposition of PM<sub>10</sub> to urban street tree foliage has been measured as 205 mg m<sup>-2</sup> (±145 mg m<sup>-2</sup>) and 361 mg m<sup>-2</sup> (±318 mg m<sup>-2</sup>) (Broadmeadow, Beckett, Jackson, Freer-Smith, & Taylor, G., 1998; Beckett, Freer-Smith, & Taylor, 2000a, 2000b). Broadmeadow et al. (1998) also reports a maximum of 1400 mg m<sup>-2</sup> of PM<sub>10</sub> deposited to foliage of an urban woodland situated 25 m from a major UK highway (the M6). These were separate studies and issues with data comparisons are given in Section 4.

With no additional expansion to the urban canopy, by 2050 the PM<sub>10</sub> removal is reduced to a total annual capture of between 693 tonnes (454 tonnes to foliage) and 1723 tonnes (1485 tonnes to foliage) for each respective model equating to a 19% reduction. This reflects the predicted decline in PM<sub>10</sub> exposure by 2050. Implementing a 30% canopy cover under scenario 1 (S1) would result in an increased annual capture of 18% above 2006 levels for both models. Under scenario 2 planting dedicated to increasing the street tree canopy (S2b) offers a slight advantage in terms of increased deposition over the other canopy types: 1109 tonnes (728 tonnes to foliage) for the pollution flux model and 2760 tonnes (2379 tonnes to foliage) for the Tiwary method. This reflects the position of street trees on roadsides where PM<sub>10</sub> ambient concentrations are greater. The pollution flux model estimates percentage reductions of PM<sub>10</sub> concentrations in the turbulent boundary layer over the whole GLA of 0.7% currently and of up to 1.1% for scenario S2b and the Tiwary method estimates a 1.4% currently and up to 2.6% for scenario S2b.

Taking into account the heterogeneous nature of PM<sub>10</sub> distribution and the greater concentrations around highways (Fig. 2) a


relationship between PM<sub>10</sub> concentration and canopy uptake was calculated (Fig. 4). The PM<sub>10</sub> concentrations used (10–80 µg m<sup>-3</sup>) are typical for the GLA which has a current mean street exposure of 23.5 µg m<sup>-3</sup> and the current maximum street tree exposure concentration was 83.4 µg m<sup>-3</sup> (LAEI, 2006). For a 96:4 broadleaf to coniferous mix and at a maximum PM<sub>10</sub> concentration (80 µg m<sup>-3</sup>), the deposition estimate from the Tiwary method (215.7 kg ha<sup>-1</sup> y<sup>-1</sup>) is double the estimate made by the pollution flux model (100.6 kg ha<sup>-1</sup> y<sup>-1</sup>). For both models a coniferous canopy sited in the most polluted areas resulted in the greatest capture of PM<sub>10</sub> ranging from 161.5 to 1653.8 kg ha<sup>-1</sup> y<sup>-1</sup> (Fig. 4b). Based on the pollution flux model the percentage reduction of PM<sub>10</sub> in the turbulent boundary layer over one hectare of canopy was 3.3% and 5.4% for broadleaved and coniferous canopies respectively and the Tiwary method estimates 5.2–54.9% reductions for broadleaved and coniferous canopies.

#### 4. Discussion

The quantification of urban canopy cover reported here (Table 1) was in agreement with those reported from other studies. The calculated average canopy cover of a garden within the GLA is within the range of cover estimated for five other UK cities (0–30%) (Loram, Warren, & Gaston, 2008).

Both the modelling approaches used here for estimation of dry deposition of PM<sub>10</sub> to the urban tree canopy estimate deposition as the inverse of three resistance parameters ( $R_a$ ,  $R_b$  and  $R_c$ ) (Baldocchi et al., 1987). When data were converted from hourly to seasonal input data those parameters computed using hourly wind speed were the most affected, for example  $R_b$  (s m<sup>-1</sup>; quasi-laminar boundary layer) for both models and the deposition velocity term ( $V_d$ ; m s<sup>-1</sup>) and  $R_c$  (canopy resistance) for the Tiwary method (Table 2). The change from using hourly to annually derived inputs had little effect on the estimates made using the pollution flux model. This suggests that the  $R_b$  parameter has little influence and that seasonally corrected annual input data can be used to estimate dry deposition of PM<sub>10</sub> to the urban canopy using the pollution flux model and that future predictions can be made from this approach. However, the estimated deposition by the Tiwary method was reduced by 30% when seasonal inputs were used. The  $V_d$  term in this approach was modelled as a near linear function with wind speed (Tiwary et al., 2009) from data collected in wind tunnels (Freer-Smith et al., 2004). This function results in very high deposition during wind gusts, for example the peak hourly wind speed in 2004 was 16.5 m s<sup>-1</sup> giving  $V_d$  values of 0.0147 m s<sup>-1</sup>





**Fig. 4.** The relationship between atmospheric  $PM_{10}$  concentrations and deposition of  $PM_{10}$  to (a) one hectare of urban canopy consisting of 96% broadleaf and 4% coniferous cover as estimated by both modelling approaches and (b) one hectare of either 100% deciduous or 100% coniferous canopies as estimated by both the pollution/deposition and Tiwary approaches. Values are given for direct model outputs; therefore values are for deposition to foliage and stems for the pollution flux and foliage alone for the Tiwary approach.

and  $1.156\ ms^{-1}$  and deposition to the whole GLA woodland of  $591\ kg\ h^{-1}$  and  $3800\ kg\ h^{-1}$  to deciduous and coniferous canopies respectively. Using the hourly input data this modelled function has been extrapolated outside of its range and the results are therefore uncertain. It has been suggested that deposition is reduced in high wind speed as leaves align themselves with wind to reduce drag, and particle bounce off becomes more common, and data from Freer-Smith et al. (2004) suggest that this near linear relationship does not exist in wind speeds greater than  $9\ ms^{-1}$ . Thus these high deposition rates estimated above the maximum wind speed from which the relationships were generated may be unrealistic. The mean seasonal wind speed for 2006 ( $3.8\ ms^{-1}$ ) was in the measured range used by Freer-Smith et al. (2004) at 4.1, 4.0, 3.3 and  $3.9\ ms^{-1}$  for winter, spring, summer and autumn respectively.

The pollution flux method gives an uptake of  $27.3\ kg\ ha^{-1}\ yr^{-1}$  to the urban canopy for the mean background  $PM_{10}$  concentrations of the GLA ( $21.68\ \mu g\ m^{-3}$  in 2006) (Table 5). This compares well with the uptake for Chicago's canopy estimated using UFORE and sub-sampling and hourly input data. Deposition to Chicago was  $30.7\ kg\ ha^{-1}\ yr^{-1}$  at annual mean  $PM_{10}$  concentrations of  $34\ \mu g\ m^{-3}$  (Nowak et al., 2006). The Tiwary method estimates  $58.5\ kg\ ha^{-1}\ yr^{-1}$  an uptake of approximately double the pollution flux model (Table 5). Both these values are within the range ( $11\text{--}80\ kg\ ha^{-1}$ ) given for annual  $PM_{10}$  removal by trees and shrubs for 55 USA cities (Nowak et al., 2006). The most recent estimates of canopy cover and the annual map of  $PM_{10}$  concentrations for 2006 gives the total deposition of  $PM_{10}$  as 852 tonnes for the pollution flux model (Fig. 4) similar to the 855 tonnes estimated by Powe and Willis (2002). Unsurprisingly, the estimated future reductions in atmospheric  $PM_{10}$  pollution (Williams, 2007) results in a reduced capture rate by the urban canopy.

For both models increasing the proportion of street trees offers the greatest potential for atmospheric  $PM_{10}$  mitigation due to the more concentrated pollutant exposure at roadsides. However, Litschke and Kuttler (2008) caution that a dense street tree canopy may not allow street derived emissions to mix with the surrounding atmosphere and therefore creating localised high  $PM_{10}$  concentrations at street level. Nevertheless, the data reported here suggest that extending the urban canopy in areas of high  $PM_{10}$  pollution offers the greatest potential for removal of  $PM_{10}$  from the atmosphere. Planting design may allow the possible decrease of dispersion ('canyon') effect to be minimized by avoiding long unbroken lines of trees or preventing the crown from closing over the top of roads. At exposures of  $80\ \mu g\ m^{-3}$  of  $PM_{10}$  the various planting scenarios reported in Fig. 4 offer annual reductions in the range 3.3–55% in  $PM_{10}$  concentrations within a well mixed boundary layer. Bealey et al. (2007) estimate that removal rates for  $PM_{10}$  of between 1% and 30% can be achieved in Glasgow depending on planting density and location but these authors also acknowledge the existence of land area constraints to planting at the most polluted sites.

Currently, in the absence of *in situ* validation for this study an attempt has been made to compare these modelled results of dry deposition with measured data of  $PM_{10}$  load to urban leaves in the UK environment. However, modelled data estimate the annual dry deposition assuming no rainfall (maximum potential dry deposition) while these measured data will have been subjected to prior rain events, and Beckett (2000a, 2000b) suggests that there is the potential for some wash off of  $PM_{10}$  by rainfall. Broadmeadow et al. (1998) supports this measuring deposition (on a leaf area basis) to an urban woodland later in the season to be less than early season values, suggesting that deposited mass was lost as the season progressed. Thus estimates of  $PM_{10}$  deposition based on leaf

washing are likely to be a useful approach to the validation of model estimates although the leaf washing potential should be considered. There is clearly a wide range in measured values of PM<sub>10</sub> deposition to urban trees. This wide range is accounted for by factors such as species differences, duration and intensity of rainfall, other meteorological factors such as wind speed, local PM<sub>10</sub> concentration, re-suspension rate and techniques for measuring PM<sub>10</sub> load and LAI.

## 5. Conclusion

Converting from hourly to seasonal input data had little impact on estimates of annual dry deposition of PM<sub>10</sub> to the urban canopy suggesting annual estimates of future predictions can be used as input data. The modelling approaches used here gave a range of annual PM<sub>10</sub> deposition to the urban canopy of London, depending on approach used and on whether trees are coniferous or broadleaved. This range is within those estimates derived for other cities using the UFORE model and the same  $V_d$  value which we used here in the pollution flux model but not in the Tiwary method (Nowak, 1994a; Nowak et al., 2006). The values derived from both the approaches described here are also very similar to those reported by Powe and Willis (2002) for London (852 tonnes and 855 tonnes respectively). Data reported here suggest that expanding the urban canopy will have a positive impact on the urban atmosphere through reduction of PM<sub>10</sub> and that coniferous tree planting in areas of higher pollution offers the greatest PM<sub>10</sub> mitigation potential. There are maintenance and opportunity costs associated with urban greenspace and when considering beneficial effects on air quality the, very species specific, potential of plants to produce biogenic volatile organic compounds (BVOCs) and their role in photochemical ozone formation also needs to be considered (see Donovan et al., 2005). There is a clear need for *in situ* validation of these modelled estimates to better parameterise the models, particularly the deposition velocity term, and improve the accuracy of modelled estimates. Sensitivity of selected species to atmospheric pollution and climate change, aesthetic appeal, biodiversity, soil factors, maintenance costs and the land availability for planting will also determine species choice and the use of trees in urban areas in future.

## Acknowledgments

This work was funded through the BRIDGE project (sustainable Urban planning Decision support accounting for urban metabolism) supported by the EC through its seventh Framework program Theme 6 contract 211345. The British Atmospheric Data Centre (BADC) is thanked for providing the hourly 2004 meteorological data. Matthew J Aylott is thanked for help and advice with ERSI® ArcMap and Dave Nowak is thanked for advice concerning UFORE.

## References

- Air Quality Expert Group. (2005). *Particulate matter in the UK: Summary*. London: Defra.
- Baldocchi, D. D., Hicks, B. B., & Camara, P. (1987). A canopy stomatal-resistance model for gaseous deposition to vegetated surfaces. *Atmospheric Environment*, 21, 91–101.
- Bealey, W. J., McDonald, A. G., Nemitz, R., Donovan, R., Dragosits, U., Duffy, T. R., et al. (2007). Estimating the reduction of urban PM10 concentrations by trees within an environmental information system for planners. *Journal of Environment Management*, 85, 44–58.
- Beckett, K. P., Freer-Smith, P. H., & Taylor, G. (2000a). The capture of particulate pollution by trees at five contrasting urban sites. *Arbor Journal*, 24, 209–230.
- Beckett, K. P., Freer-Smith, P. H., & Taylor, G. (2000b). Effective tree species for local air-quality management. *Arbor Journal*, 26, 12–19.
- Beckett, K. P., Freer-Smith, P. H., & Taylor, G. (2000c). Particulate pollution capture by urban trees: Effect of species and windspeed. *Global Change Biology*, 6, 995–1003.
- Bower, J., Broughton, G., Connolly, C., Cook, A., Eaton, S., Glynn, A., et al. (2008). Air pollution in the UK: 2008. Crown Copyright, AEAT/ENV/R/2823/Issue 1.
- Britt, C., & Johnston, M. (2008). *Trees in towns II: A new survey of urban trees in England and their condition and management. Research for amenity trees; No. 9*. Department for Communities and Local Government.
- Broadmeadow, M., Beckett, P., Jackson, S., Freer-Smith, P., & Taylor, G. (1998). *Forest research annual report and accounts 1997–98* (pp. 35–43). TSO. ISBN: 0 10 254598 7.
- Colbeck, I., & Hanison, R. M. (1985). Dry deposition of ozone: Some measurements of deposition velocity and of vertical profiles to 100 metres. *Atmosphere Environment*, 19(11), 1807–1818.
- COMEAP. (1998). *Quantification of the effects of air pollution on health in the United Kingdom*. London: Committee of the Medical Effects of Air Pollutants, Department of Health, The Stationery Office.
- Donovan, R. G., Stewart, H. E., Owen, S. O., MacKenzie, A. R., & Hewitt, C. N. (2005). Development and application of an urban tree air quality score for photochemical pollution episodes using the Birmingham, United Kingdom, area as a case study. *Environmental Science & Technology*, 39, 6730–6738.
- Fowler, D., Cape, T. N., & Unsworth, M. H. (1989). Deposition of atmospheric pollutants on forests. *Philosophical Transactions of the Royal Society of London*, 324, 247–265.
- Fowler, D., Skiba, U., Nemitz, E., Choubedar, F., Branford, D., Donovan, R., et al. (2004). Measuring aerosol and heavy metal deposition on urban woodland and grass using inventories of 210Pb and metal concentrations in soil. *Water, Air and Soil Pollution*, 4, 483–499.
- Freer-Smith, P. H., Beckett, K. P., & Taylor, G. (2005). Deposition velocities to *Sorbus aria*, *Acer campestre*, *Populus deltoides* X *trichocarpa* 'Beaupre', *Pinus nigra* and *X Cupressocypariss leylandii* for coarse, fine and ultra-fine particles in the urban environment. *Environmental Pollution*, 133, 157–167.
- Freer-Smith, P. H., El-Khatib, A. A., & Taylor, G. (2004). Capture of particulate pollution by trees: A comparison of species typical of semi-arid areas (*Ficus nitida* and *Eucalyptus globulus*) with European and North American species. *Water, Air and Soil Pollution*, 155, 173–187.
- Fuller, R., Smith, G. M., Sanderson, J. M., Hill, R. A., & Thompson, A. G. (2001). The UK Land Cover Map 2000: Construction of a parcel-based vector map from satellite images. *Cartographic Journal*, 39, 15–25.
- Fuller, R., Smith, G. M., Sanderson, J. M., Hill, R. A., Thompson, A. G., Cox, R., et al. (2002). Countryside Survey 2000 Module 7. Land cover map 2000 a guide to the classification system. Extract from the draft final report cslcm/final. Centre for Ecology and Hydrology (Natural Environment Research Council) Project. T02083j5/C00878.
- GLA. (2007). *Chainsaw massacre: A review of London's street trees*. Greater London Authority.
- GLA. (2009). *Leading to a greener London: An environment programme for the capital*. Greater London Authority.
- Gupta, A., Kumar, R., Maharaj Kumari, K., & Srivastava, S. S. (2004). Atmospheric dry deposition to leaf surfaces at a rural site of India. *Chemosphere*, 55, 1097–1107.
- LAEI. (2006). London Atmospheric Emissions Inventory 2006. Compiled by AEA, March 2009. Available from [http://static.london.gov.uk/mayor/environment/air\\_quality/research/emissions-inventory.jsp](http://static.london.gov.uk/mayor/environment/air_quality/research/emissions-inventory.jsp).
- Litschke, T., & Kuttler, W. (2008). On the reduction of urban particle concentration by vegetation – A review. *Meteorologische Zeitschrift*, 17, 229–240.
- Loram, A., Warren, P. H., & Gaston, K. J. (2008). Urban domestic gardens (XIV): The characteristics of gardens in five cities. *Environmental Management*, 42, 361–376.
- Lovett, G. M. (1994). Atmospheric deposition of nutrients and pollutants in North America: An ecological perspective. *Ecological Applications*, 4, 629–650.
- Meteorological Office. (2006). Meteorological Office Integrated Data Archive System (MIDAS) land surface stations data (1853–current) [Internet]. British Atmospheric Data Centre. Available from <http://badc.nerc.ac.uk/data/ukmo-midas>.
- Nowak, D. J. (1994a). Air pollution removal by Chicago's urban forest. In E. G. McPherson, D. J. Nowak, & R. A. Rowntree (Eds.), *Chicago's urban forest ecosystem: Results of the Chicago urban forest climate project* (pp. 63–82). Radnor, PA: US Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
- Nowak, D. J. (1994b). Urban forest structure: The state of Chicago's urban forest. In E. G. McPherson, D. J. Nowak, & R. A. Rowntree (Eds.), *Chicago's urban forest ecosystem: Results of the Chicago urban forest climate project* (pp. 3–18). Radnor, PA: US Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
- Nowak, D. J., & Crane, D. E. (1998). The Urban Forest Effects (UFORE) model: Quantifying urban forest structure and functions. In *Integrated tools proceedings*.
- Nowak, D. J., Crane, D. E., & Stevens, J. C. (2006). 2006 air pollution removal by urban trees and shrubs in the United States. *Urban Forestry and Urban Greening*, 4, 115–123.
- Perry, M., & Hollis, D. (2005). The generation of monthly gridded datasets for a range of climatic variables over the UK. *International Journal of Climatology*, 25(8), 1041–1054. Current spatial maps of met data from: <http://www.metoffice.gov.uk/climatechange/science/monitoring/ukcp09/download/monthly/gridded-monthly.html>
- Powe, N. A., & Willis, K. G. (2002). Social and environmental benefits of forestry. Phase 2: Mortality and morbidity benefits of air pollution absorption by woodland. Report to Forestry Commission.
- Powe, N. A., & Willis, K. G. (2004). Mortality and morbidity benefits of air pollution (SO<sub>2</sub> and PM<sub>10</sub>) absorption attributable to woodland in Britain. *Journal of Environment Management*, 70, 119–128.
- Read, D. J., Freer-Smith, P. H., Morison, J. I. L., Hanley, N., West, C. C., & Snowdon, P. (Eds.). (2009). *Combating climate change – A role for UK forests: An assessment*

- of the potential of the UK's trees and woodlands to mitigate and adapt to climate change*. Edinburgh: The Stationary Office.
- Smith, W. H. (1981). *Air pollution and forests*. NY: Springer-Verlag.
- Tiwary, A., Sinnett, D., Peachy, C., Chalabi, Z., Vurdoulakis, S., Fletcher, T., et al. (2009). An integrated tool to assess the role of new planting in PM10 capture and the human health benefits: A case study in London. *Environmental Pollution*, 157, 2645–2653.
- UKCP. (2009). UK climate projections 2009. Available from <http://ukclimateprojections.defra.gov.uk/>.
- UKCP02 Hulme, M., Jenkins, G. J., Lu, X., Turnpenny, J. R., Mitchell, T. D., Jones, R. G., et al. (2002). *Climate change scenarios for the United Kingdom: The UKCIP02 scientific report*. Norwich, UK: Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia.
- Williams, M. L. (2007). UK air quality in 2050 – Synergies with climate change policies. *Environmental Science Policy*, 10, 169–175.
- Yang, J., McBride, J., Zhou, J., & Sun, Z. (2005). The urban forest in Beijing and its role in air pollution reduction. *Urban Forestry and Urban Greening*, 3, 65–78.
- Zinke, P. J. (1967). Forest interception studies in the United States. In W. E. Sopper, & H. W. Lull (Eds.), *Forest hydro* (pp. 137–161). Oxford: Pergamon Press.